

**SURGICAL ASSOCIATES, PC**  
**GENERAL, THORACIC AND LAPAROSCOPIC SURGERY**  
2900 12<sup>TH</sup> AVE NORTH \* SUITE 355W  
BILLINGS, MT 59101-0137  
(406) 238-6470 \* (406) 238-6499

DENNIS W. MAIER, M.D., F.A.C.S.  
GEORGE K. BENTZEL, M.D., F.A.C.S.  
JEFFREY J. RENTZ, M.D., F.A.C.S.  
BARRY A. MCKENZIE, M.D., F.A.C.S.

ERIC R. DRINGMAN, M.D., F.A.C.S.  
KATHRYN F. HATCH, M.D., F.A.C.S.  
MICHAEL G. WILCOX, M.D., F.A.C.S.

**AFTER CARE**  
**HEMORRHOID SURGERY**

I. **COMMON FINDINGS INCLUDE:**

**Pain.** It is normal to pass blood clots and small amounts of bright red blood for several days.

II. **CALL YOUR SURGEON FOR THE FOLLOWING:**

- (1) Sustained fever >100 degrees
- (2) Significant or prolonged bleeding

III. **COMMON SIDE EFFECTS OF PAIN MEDICATIONS INCLUDE:**

- (1) Nausea – Be sure to take medication with some food or drinks.
- (2) Constipation –See bowel regimen below
- (3) Light-headedness
- (4) Do not drive while taking narcotic pain medication

If these symptoms occur after taking medications, discontinue use.  
May take over the counter pain medications or laxative of choice.

IV. **BOWEL REGIMEN TO AVOID CONSTIPATION:**

- (1) Colace 100mg to keep stools soft. Reduce for watery or loose stools. Stop after one week or after stopping pain medications.
- (2) Milk of Magnesia. You should have 1 bowel movement daily. If not, then take Milk of Magnesia the next morning. Drink Plenty of water. Recommend drinking eight 8oz glasses of water daily.
- (3) Fiber supplements. Take prescribed amount of fiber in double water.
- (4) May drink prune juice.

**NOTE:** Routine calls and prescription refills are best handled during office hours 8am-5pm Monday through Friday. For urgent problems after hours, the surgeon on call can be reached at (406) 238-6470.